



*A sandstorm hovering over the African Union-United Nations Mission in North Darfur, El Fasher, Sudan
Photo credit: UNAMID/Adrian Dragnea*

Sand and Dust Storms: Subduing a Global Phenomenon

Invasion by sand and dust

In 2010, Chinese authorities issued a level-five pollution warning as a massive sandstorm moved from Mongolia and northern China towards Beijing, hovering over an area of 810 000 square kilometres and threatening 250 million people.¹ In May 2016, a series of massive sand storms swept across Rigan County in southeast Iran, burying 16 villages and causing USD 9 million in losses.² A few months later, heavy dust and sand clouds engulfed Abu Dhabi, reducing the visibility in the city to 500 metres and increasing the number of asthma patients admitted to hospitals by 20 per cent.^{3,4} These are just a few examples of recent threats and damages that sand and dust storms delivered to many parts of the world. Other examples abound throughout human history.⁵

Sand and dust storms result when strong turbulent winds erode sand and silt particles from arid and semi-arid landscapes and then launch them into the air. Sand storms move relatively close to the ground and both particle size and wind speed constrain the distance each particle travels. Dust storms raise large quantities of fine silt particles and smaller clay particles higher into the atmosphere.⁶

Dust storms can travel thousands of kilometres across continents and oceans, entraining other pollutants on the way and depositing particles far from their origin. Winds spread the dust of the Sahara—the most significant source—west to the Americas, north to Europe and east to China.⁶ Central Asian and Chinese sources reach the Korea peninsula, Japan, the Pacific Islands, North America and beyond.




A 2003 case study traced large amounts of dust from a 1990 Chinese plume that reached the European Alps—travelling eastward more than 20 000 kilometres within two weeks.⁷ Dust plays an important role in biogeochemical processes throughout the Earth system. It forms a source material for vast expanses of loess soils.⁸ Mineral dust deposition provides nutrients such as iron and other trace elements to terrestrial and marine ecosystems, boosting primary productivity and phytoplankton growth.⁹ Saharan dust is a natural fertilizer of the Amazon rainforest, providing phosphorus inputs that balance losses through river discharge.¹⁰ Similarly, Hawaiian rain forests receive nutrients from the dust of Central Asia.¹¹ At the same time, dust from Africa and Asia may damage Caribbean coral reefs.¹²

Dust can also harm animals and humans, especially in arid and semi-arid regions. For humans, inhaling fine particles can generate and aggravate asthma, bronchitis, emphysema, and

silicosis.¹³ Finer dust also can deliver a range of pollutants, spores, bacteria, fungi and allergens. Other common problems include eye infections, skin irritations and Valley Fever. In countries of the Sahel, dust loads arriving from the Sahara correlate strongly with meningitis outbreaks.¹⁴ Chronic exposure to fine dust contributes to premature death from respiratory and cardiovascular diseases, lung cancer and acute lower respiratory infections.¹⁵

Other socioeconomic damages follow dust storms.¹⁶⁻¹⁹ Short-term costs include disease and death in livestock, crop destruction, damage to buildings and other infrastructure, transport shutdowns and expensive removal of tonnes of deposits. Economic losses from a single storm event can be hundreds of millions of dollars. Longer-term costs include erosion of soils, pollution of ecosystems, chronic debilitating health problems, and desertification.



A portrait of global aerosols produced by a GEOS-5 simulation at a 10-kilometre resolution. Dust emission can be seen in brown/red colour.

Photo credit: William Putman, NASA/Goddard Space Flight Center

Drivers from nature, land mismanagement, and climate change


Dust activities vary significantly on a variety of time scales, such as seasonal, annual, decadal or multi-decadal scales.²⁰ A 2012 study of satellite data from 2003 to 2009 in comparison with similar analyses of data from earlier periods suggests that over the past three decades, substantial changes have occurred in Australia, Central Asia and the US high plains; whereas dust events over Northern Africa, the Middle East and South America have remained at the same level of activity.^{21,22} Further studies show that these regions experience frequent high dust intensity as storms or haze events derived from both natural and anthropogenic causes.²¹⁻²³

The anthropogenic causes, responsible for about 25 per cent of global dust emission, result from land use changes that include excessive water extraction and water diversion for irrigation, leading to drying out of water bodies; and deforestation and unsustainable agricultural practices, which expose soils to wind erosion. These are all forms of land degradation. In drylands, when agricultural soils are tilled too often and too deeply and crop residues are removed, soils are left exposed. Removal of hedges and windrows to allow for larger equipment increases wind erosion. Overgrazing of rangelands results in loss of soil cover. When the soils have no ground cover, winds lift away the finer particles containing much of the soil's nutrients and organic matter. Model simulations suggest that globally, dust emissions have increased between 25 to 50 per cent due to a combination of land use and climate change since 1900.²⁴

In every dust-prone region, the relationship between human activities and increased dust is visible and sometimes palpable. The Owens (dry) Lake dust source in California was desiccated after water diversions to the Los Angeles Aqueduct began in 1913.²⁵ Patagonia in the southern half of Argentina has become a major anthropogenic dust source from desertification caused by unsustainable ranching.²⁶ The Indo-Gangetic basin is a principal dust source in South Asia, resulting from intensive agricultural activities.²² In Australia, land clearing and water demand for agriculture has disrupted the hydrological regime and led to significant increase in dust.²⁷ Lake Balkhash in Kazakhstan has been rapidly drying since 1970 after completion of a dam upstream on the Ili River.

Shrinking Aral Sea from 2000 to 2013

Following decades of large-scale water diversion, the Aral Sea has dried up and become active sources of dust



Credits

1989 - University of Maryland Global Land Cover Facility

2003 - Jacques Descloitres, NASA/Goddard Space Flight Center

2014 - Jesse Allen, NASA Earth Observatory

And finally, decades of large-scale diversions—from the region's main rivers, the Syr Darya and Amu Darya, to extensive irrigation schemes—reduced river flow reaching the Aral Sea, with consequent desiccation and desertification throughout the region.²⁸ Vast areas of the Aral Sea Basin are now active sources of noxious dust polluted with the persistent residue of artificial fertilizers and pesticides that were banned from use decades ago.²⁹

Anthropogenic climate change is an important driver of dust generation, in addition to that produced naturally and by unwise land management. Many regions that are currently dusty areas will likely become drier and contribute more atmospheric dust. These include most of the Mediterranean areas of Africa and Europe, northern Sahara, West Asia, Central Asia, southwest USA, and southern Australia.^{30,31} In turn, increased dust in the atmosphere can influence the climate system. It can disturb the earth's radiative balance, intensifying droughts in arid areas.³² On the other hand, dust could enhance precipitation in some areas, by seeding clouds.³³

Sand and dust storms are thus inter-connected with a range of environmental and development issues that extend across national, regional, and continental bounds. Anthropogenic climate change will further exacerbate decades of unsustainable land and water resource management in regions that generate sand and dust storms. This threat can be diminished by quick and effective action.

Sources and impacts of sand and dust storms

Land use changes such as agriculture, water diversion and deforestation generate **25% of global dust**

Since 1900 dust emissions have **increased by 25-50%** due to human activity

Sand and dust storms are common in **arid** and **semi-arid** areas

As **climate changes**, more variability and extreme events increase the risk from dust storms

Ecological restoration initiatives help reduce the frequency and severity of local dust storms

Sand and dust storms contain **particles** in a **wide range of sizes**

Sand and dust storms occur when strong turbulent winds erode and lift particles of sand and silt from arid landscapes

When inhaled, particles smaller than **10 microns**—one-hundredth of a millimetre—cause heart and lung disease

In dust storms, dust concentrations reach **100-1 000 $\mu\text{g}/\text{m}^3$**

During an Iranian dust storm in January 2017, fine particle concentrations exceeded **10 000 $\mu\text{g}/\text{m}^3$**

WHO sets safe air quality for concentration of fine particles at or below **50 $\mu\text{g}/\text{m}^3$**


Dust storms can carry a range of **pollutants, spores, fungi, bacteria and allergens**. Dust carried from the Sahara desert can cause Meningitis outbreaks in the Sahel.

A dust storm in northwest China in 1993 killed nearly **120 000 livestock**; destroyed **373 333 hectares of crops**; and buried over 2 000 km of irrigation ditches

Economic losses from a single dust storm can reach **hundreds of millions of dollars**

Dust storms damage crops, kill livestock and erode fertile soil

Arid regions are likely to become drier and face more dust storms, including the **Mediterranean** areas of **Europe** and **Africa**, **northern Sahara**, **central and west Asia**, **southwest USA**, and **southern Australia**



Reducing damage through focus on smaller scales

In the short to medium-term, successful efforts to reduce threats from sand and dust storms should focus on protective strategies.³⁴ Of course, early warning systems and disaster reduction procedures are essential components of preparedness, and regional programmes are evolving to refine these services. Procedures for coping in real time with sand and dust storms include advisory communications to public services; school, airport, rail and road closures; and hospital emergency services.

Preparedness starts with public awareness of sand and dust storm risks through education in schools, media and social networks and telecommunication. Preparedness should also include techniques that provide physical protection of valuable assets such as planting or erecting barriers upwind of populated areas and essential infrastructure to encourage dust deposition outside those areas. Some actions, such as aligning roads and removing traps, channel dominant winds and their burden away from sites needing protection.

In the medium to long-term, reducing the threat from sand and dust storms should focus on preventive strategies


Video: Africa's ambitious "Great Green Wall"



Video Link: https://www.youtube.com/watch?v=jL_nRHg-0l4

Photo Credit: Tilling soil in Senegal by IFPRI/Milo Mitchell, licensed under CC BY-NC-ND 2.0

© TIME



A dust plume blowing from Northern Africa towards Europe and the Atlantic Ocean, 21 February 2017


Photo credit: NASA image by Jeff Schmaltz, LANCE/EOSDIS Rapid Response

that promote sustainable land and water management across landscapes. This level of scope addresses cropland, rangelands, deserts, and urban areas. Such strategies should integrate with measures for climate change adaptation and mitigation, as well as for conserving biological diversity. These crucial integrated strategies are currently deficient in many vulnerable regions.³⁴

The Three North Shelterbelt Program in China, sometimes called the Green Great Wall, is an integrated effort that began in 1978 to address rampant soil erosion, which exacerbated existing problems with flooding and dust storms over large expanses, after decades of unsustainable natural resource exploitation. Research results and lessons learned suggest that a focus on what works at community and local levels, with local plant species already adapted to specific locations, brings success when linked and scaled up.³⁵ These insights renew emphasis on actions that promote ecosystem services such as food production, carbon sequestration, soil and water retention, flood mitigation, and provision of habitat for biodiversity conserving natural capital, as well as prevention of sand and dust storms.³⁶ Observations on Green Great Wall efforts show significant improvements in the surrounding vegetation index and deduce these efforts have effectively reduced dust storm intensity, after accounting for influences from climatic change and human pressures.^{37,38}



Sources of dust emissions (tonnes/year)




Data source: Ginoux et al. (2012)³²

In the Kubuqi desert of Inner Mongolia, private-public-community investments to plant local species of trees, shrubs and grasses on more than 5 000 square kilometres of desert land reduced the frequency of dust storms and the amount of related damage to homes and infrastructure.³⁹

In Africa, the Great Green Wall for the Sahara and the Sahel Initiative is also finding success by working at local and community scales.⁴⁰ The initiative has graduated from a tree planting vision to focus on broader sustainable development: In Senegal they started by planting more than 270 square kilometres of indigenous trees that do not need watering. Later, other plants and animals are reappearing to restore the ecosystem. Communities in Mauritania, Chad, Niger, Ethiopia and Nigeria are growing market gardens along the dryland edge giving the young population work and reasons to reject migration. Once again, to succeed these projects carefully select plant species that are well adapted to local conditions, matched to water resource availability, and familiar to local residents who will ultimately be responsible for maintaining the restoration and the landscape.⁴¹

Economic losses due to sand and dust storms



Multilateral support for reducing sand and dust storm damage

Integrated strategies that address sand and dust storm threats reflect recommended actions to contain land degradation, terrestrial biodiversity loss, and climate change threats under the three Rio Conventions: the UN Convention to Combat Desertification (UNCCD), the UN Convention on Biological Diversity and the UN Framework Convention on Climate Change, respectively. Supported by the UNCCD, West Asia and Northeast Asia have developed Regional Action Plans on sand and dust storms and the Northeast Asia plan is in full operation.⁴²

Each Rio Convention supports land and water management efforts in partnership with appropriate multilateral institutions and agencies. International unity on these issues is reflected in the Sustainable Development Goals—particularly Goal 1, 2, 5, 13 and 15—that address the integrity and management of land and water resources, and specifically Target 15.3: “By 2030, combat desertification, restore degraded land and soil, including land affected by desertification, drought and floods, and strive to achieve a land degradation-neutral world.”


 **Video: Combatting desertification: Chinese herdsmen dedicated to turn desert into oasis**



Video Link: <https://www.youtube.com/watch?v=gITXPUrYYJ0>

© CCTV English

Photo credit: Preventing desertification in Ningxia, China by Bert van Dijk, licensed under CC BY-NC-SA 2.0



A dust storm over the Persian Gulf, 19 February 2017

Photo credit: NASA image by Jeff Schmaltz, LANCE/EOSDIS Rapid Response

Regional frameworks, agreements and action plans such as the Regional Master Plan for the Prevention and Control of Dust and Sand Storms in Northeast Asia and national action plans, such as those required by the UNCCD, also establish policies for reducing sand and dust storm threats.


The World Meteorological Organization has established the Sand and Dust Storm Warning Advisory and Assessment System to enhance the ability of countries to deliver early and accurate sand and dust storm forecasts, observations, information and knowledge to users.⁴³ This system provides global and regional forecasts of dust threat and has established regional centres for the Americas, Asia, and Northern Africa, Middle East and Europe.⁴⁴

Integrated land and water management efforts encompass agricultural activities, and the Food and Agriculture Organization promotes conservation agriculture to address threats in arid regions. In 1992, a network called the World Overview of Conservation Approaches and Technologies (WOCAT) began to collect information about conservation




Online dust forecast by the World Meteorological Organization Sand and Dust Storm Warning, Advisory and Assessment System Regional Centres


Dust Surface Concentration (ug/m³) 18 May 2017



Surface Dust Concentration (ug/m³) 18 May 2017



Surface Dust Concentration 21 May 2017



agriculture and sustainable land management practices from specialists. In 2014 the network was formalized into a consortium and recognized by the UNCCD as the recommended source of data on best practice. In 2017, WOCAT involves more than 2 000 registered users, over 60 participating institutions, and around 30 national and regional initiatives.⁴⁵

Agriculture is responsible for nearly 70 per cent of all freshwater withdrawals.⁴⁶ Conservation agriculture also promotes water use practices that prevent water scarcity and desertification and that reduce the formation of sand and dust storms. The 2030 Water Resources Group has brought together case studies from around the world of currently available, replicable and practical solutions for water management. The solutions have been collected in an online catalogue 'Managing Water Use in Scarce Environments' that is meant to inspire action and use by policy and decision makers.⁴⁷ Many of the solutions have clear relevance to reducing sand and dust storms.

Finally, improved international integration and coordination of research is needed to reduce critical uncertainties on the interaction of dust with global biogeochemical processes and climate systems; to improve methods for monitoring, prediction and early warning systems; to assess the economic impacts and costs of sand and dust storms and related mitigation measures; and to improve the effectiveness of measures before, during and after interventions.

1 Northern Africa, Middle east and Europe Centre
<https://sds-was.aemet.es/>

2 WMO SDS-WAS Asian Center
http://eng.nmc.cn/sds_was.asian_rc/

3 Pan-America Regional Center
<http://sds-was.cimh.edu.bb/>

References

1. BBC (2010). China sandstorm leaves Beijing shrouded in orange dust. *BBC*, 20 March 2010. <http://news.bbc.co.uk/2/hi/asia-pacific/8577806.stm>
2. Tehran Times (2016). Sand storm buries 16 villages in southeastern Iran. *Tehran Times*, 18 May 2016. <http://www.tehrantimes.com/news/402617/Sand-storm-buries-16-villages-in-southeastern-Iran>
3. Emirates 24/7 News (2016). NCMS warns of active winds, low visibility. *Emirates 24/7 News*, 4 August 2016. <http://www.emirates247.com/news/emirates/ncms-warns-of-active-winds-low-visibility-2016-08-04-1.637979>
4. The National (2016). Asthma attacks on the rise in UAE as winds whip up sand and dust. *The National*, 19 July 2016. <http://inbusiness.ae/2016/07/19/asthma-attacks-on-the-rise-in-uae-as-winds-whip-up-sand-and-dust>
5. McLeman, R., Dupre, J., Berrang Ford, L., Ford, J., Gajewski, K. and Marchildon, G. (2014). What We Learned from the Dust Bowl: Lessons in Science, Policy, and Adaptation. *Population and Environment*, 35, 417–440. <https://www.ncbi.nlm.nih.gov/pubmed/24829518>
6. Goudie, A.S. and Middleton, N.J. (2006). *Desert Dust in the Global System*. Springer-Verlag Berlin Heidelberg
7. Grousset, F.E., Ginoux, P. and Bory, A. (2003). Case study of a Chinese dust plume reaching the French Alps. *Geophysical Research Letters*, 30(6), 1277. <http://onlinelibrary.wiley.com/doi/10.1029/2002GL016833/full>
8. Pye, K. (1987). *Aeolian dust and dust deposits*. Academic Press, London
9. Wang, F., Zhao, X., Gerlein-Safdi, C., Mu, Y., Wang, D. and Lu, Q. (2017). Global sources, emissions, transport and deposition of dust and sand and their effects on the climate and environment: a review. *Frontiers of Environmental Science & Engineering*, 11, 13. <https://link.springer.com/article/10.1007/s11783-017-0904-z>
10. Yu, H., Chin, M., Yuan, T., Bian, H., Remer, L.A., Prospero, J.M., Omar, A., Winker, D., Yang, Y., Zhang, Y., Zhang, Z. and Zhao, C. (2015). The fertilizing role of African dust in the Amazon rainforest: A first multiyear assessment based on data from Cloud–Aerosol Lidar and Infrared Pathfinder Satellite Observations. *Geophysical Research Letters*, 42, 1984–1991. https://www.researchgate.net/publication/272754426_The_Fertilizing_Role_of_African_Dust_in_the_Amazon_Rainforest_A_First_Multiyear_Assessment_Based_on_CALIPSO_Lidar_Observations
11. Chadwick, O.A., Derry, L.A., Vitousek, P.M., Huebert, B.J. and Hedin, L.O. (1999). Changing sources of nutrients during four million years of ecosystem development. *Nature*, 397, 491–497. <https://www.nature.com/nature/journal/v397/n6719/pdf/397491a0.pdf>
12. Garrison, V.H., Shinn, E.A., Foreman, W.T., Griffin, D.W., Holmes, C.W., Kellogg, C.A., Majewski, M.S., Richardson, L.L., Ritchie, K.B. and Smith, G.W. (2003). African and Asian dust: from desert soils to coral reefs. *BioScience*, 53, 469–480. <https://academic.oup.com/bioscience/article/53/5/469/241414/African-and-Asian-Dust-From-Desert-Soils-to-Coral>
13. Derbyshire, E. (2007). Natural minerogenic dust and human health. *AMBIO: A Journal of the Human Environment*, 36, 73–77. https://www.wou.edu/las/physci/taylor/g473/med_geo/derbyshire_2007.pdf
14. García-Pando, C.P., Stanton, M.C., Diggle, P.J., Trzaska, S., Miller, R.L., Perlwitz, J.P., Baldasano, J.M., Cuevas, E., Ceccato, P., Yaka, P. and Thomson, M.C. (2014). Soil dust aerosols and wind as predictors of seasonal meningitis incidence in Niger. *Environmental Health Perspectives*, 122(7), 679–686. <https://ehp.niehs.nih.gov/wp-content/uploads/122/7/ehp.1306640.pdf>
15. WHO (2013). *Review of evidence on health aspects of air pollution – REVIHAAP Project*. World Health Organization Regional Office for Europe, Copenhagen. http://www.euro.who.int/__data/assets/pdf_file/0020/182432/e96762-final.pdf
16. Tozer, P. and Leys, J. (2013). Dust storms – what do they really cost? *The Rangeland Journal*, 35, 131–142. <http://www.publish.csiro.au/rj/pdf/RJ12085>
17. Miri, A., Ahmadi, H., Ekhtesasi, M.R., Panjehkeh, N. and Ghanbari, A. (2009). Environmental and socio-economic impacts of dust storms in Sistan Region, Iran. *International Journal of Environmental Studies*, 66(3), 343–355. <http://www.tandfonline.com/doi/abs/10.1080/00207230902720170?journalCode=genv20>
18. Almasi, A., Mousavi, A.R., Bakhshi, S. and Namdari, F. (2014). Dust storms and environmental health impacts. *Journal of Middle East Applied Science and Technology*, 8, 353–356. https://www.researchgate.net/publication/271211840_Dust_storms_and_environmental_health_impacts
19. Stefanski, R. and Sivakumar, M.V.K. (2009). Impacts of Sand and Dust Storms on Agriculture and Potential Agricultural Applications of a SDSWS. *IOP Conference Series: Earth and Environmental Science*, 7(1), 012016. <http://iopscience.iop.org/article/10.1088/1755-1307/7/1/012016/pdf>
20. Shao, Y., Klose, M. and Wyrwoll, K.H. (2013). Recent global dust trend and connections to climate forcing. *Journal of Geophysical Research: Atmospheres*, 118, 1–12. https://www.researchgate.net/publication/263182073_Recent_global_dust_trend_and_connections_to_climate_forcing_GLOBAL_DUST_TREND
21. Prospero, J.M., Ginoux, P., Torres, O., Nicholson, S.E. and Gill, T.E. (2002). Environmental characterization of global sources of atmospheric soil dust identified with the Nimbus 7 Total Ozone Mapping Spectrometer (TOMS) absorbing aerosol product. *Reviews of Geophysics*, 40, 2–31. <http://onlinelibrary.wiley.com/doi/10.1029/2000RG000095/full>
22. Ginoux, P., Prospero, J.M., Gill, T.E., Hsu, N.C. and Zhao, M. (2012). Global-scale attribution of anthropogenic and natural dust sources and their emission rates based on MODIS Deep Blue aerosol products. *Reviews of Geophysics*, 50. <http://onlinelibrary.wiley.com/doi/10.1029/2012RG000388/epdf>





23. Stanelle, T., Bey, I., Raddatz, T., Reick, C. and Tegen, I. (2014). Anthropogenically induced changes in twentieth century mineral dust burden and the associated impact on radiative forcing. *Journal of Geophysical Research: Atmospheres*, 119, 526–546. <http://onlinelibrary.wiley.com/doi/10.1002/2014JD022062/epdf>
24. Mahowald, N.M., Kloster, S., Engelstaedter, S., Moore, J.K., Mukhopadhyay, S., McConnell, J.R., Albani, S., Doney, S.C., Bhattacharya, A., Curran, M.A.J. and Flanner, M.G. (2010). Observed 20th century desert dust variability: impact on climate and biogeochemistry. *Atmospheric Chemistry and Physics*, 10, 10875–10893. https://www.mpimet.mpg.de/fileadmin/staff/klosterville/Mahowald_et_al_ACPD_2010.pdf
25. Gill, T.E. (1996). Eolian sediments generated by anthropogenic disturbance of playas: Human impacts on the geomorphic system and geomorphic impacts on the human system. *Geomorphology*, 17, 207–228. https://www.researchgate.net/publication/222233193_Eolian_sediments_generated_by_anthropogenic_disturbance_of_playas_Human_impacts_on_the_geomorphic_system_and_geomorphic_impacts_on_the_human_system
26. McConnell, J.R., Aristarain, A.J., Banta, J.R., Edwards, P.R. and Simões, J.C. (2007). 20th-Century doubling in dust archived in an Antarctic Peninsula ice core parallels climate change and desertification in South America. *Proceedings of the National Academy of Sciences*, 104(14), 5743–5748. <http://www.pnas.org/content/104/14/5743.full.pdf>
27. Marx, S.K., Kamber, B.S., McGowan, H.A. and Denholm, J. (2011). Holocene dust deposition rates in Australia's Murray-Darling Basin record the interplay between aridity and the position of the mid-latitude westerlies. *Quaternary Science Reviews*, 30(23), 3290–3305. https://www.researchgate.net/publication/232391398_Holocene_dust_deposition_rates_in_Australia's_Murray-Darling_Basin_record_the_interplay_between_aridity_and_the_position_of_the_mid-latitude_westerlies
28. Groll, M., Opp, C. and Aslanov, I. (2012). Spatial and temporal distribution of the dust deposition in Central Asia – results from a long term monitoring program. *Aeolian Research*, 9, 49–62. https://www.researchgate.net/publication/257708671_Spatial_and_temporal_distribution_of_the_dust_deposition_in_Central_Asia_-_results_from_a_long_term_monitoring_program
29. Ataniyazova, O.A. (2003). *Health and ecological consequences of the Aral Sea crisis*. In the 3rd World Water Forum, Regional Cooperation in Shared Water Resources in Central Asia, Kyoto, March 18 2003, Panel III: Environmental Issues in the Aral Sea Basin. http://www.cae.utexas.edu/prof/mckinney/ce385d/papers/atanizaova_wwf3.pdf
30. Christensen, J.H., Hewitson, B., Busiuc, A., Chen, A., Gao, X., Held, I., Jones, R., Kolli, R.K., Kwon, W.-T., Laprise, R., Magaña Rueda, V., Mearns, L., Menéndez, C.G., Räisänen, J., Rinke, A., Sarr, A. and Whetton, P. (2007). Regional Climate Projections. In *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change* [Solomon, S., D. Qin, M. Manning, Z. Chen, M. Marquis, K.B. Averyt, M. Tignor and H.L. Miller (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA. <https://www.ipcc.ch/pdf/assessment-report/ar4/wg1/ar4-wg1-chapter11.pdf>
31. IPCC (2013). Summary for Policymakers. In *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Stocker, T.F., D. Qin, G.-K. Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia, V. Bex and P.M. Midgley (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA. http://www.ipcc.ch/pdf/assessment-report/ar5/wg1/WG1AR5_TS_FINAL.pdf
32. Han, Y., Dai, X., Fang, X., Chen, Y. and Kang, F. (2008). Dust aerosols: a possible accelerant for an increasingly arid climate in North China. *Journal of Arid Environments*, 72(8), 1476–1489. <http://www.sciencedirect.com/science/article/pii/S0140196308000372>
33. Twohy, C. H., Kreidenweis, S. M., Eidhammer, T., Browell, E. V., Heymsfield, A. J., Bansemer, A. R., Anderson, B. E., Chen, G., Ismail, S., DeMott, P. J. and Van den Heever, S. C. (2009). Saharan dust particles nucleate droplets in eastern Atlantic clouds. *Geophysical Research Letters*, 36, L01807. <http://onlinelibrary.wiley.com/doi/10.1029/2008GL035846/epdf>
34. UNEP, WMO and UNCCD (2016). *Global Assessment of Sand and Dust Storms*. United Nations Environment Programme, Nairobi. https://unep.unep.org/media/docs/assessments/global_assessment_of_sand_and_dust_storms.pdf
35. Xu, J. (2011). China's new forests aren't as green as they seem: impressive reports of increased forest cover mask a focus on non-native tree crops that could damage the ecosystem. *Nature*, 477(7365), 371–372. <http://www.nature.com/news/2011/110921/full/477371a.html>
36. Ouyang, Z., Zheng, H., Xiao, Y., Polasky, S., Liu, J., Xu, W., Wang, Q., Zhang, L., Xiao, Y., Rao, E. and Jiang, L. (2016). Improvements in ecosystem services from investments in natural capital. *Science*, 352(6292), 1455–1459. http://csis.msu.edu/sites/csis.msu.edu/files/Ecosystems_China_2016.pdf
37. Tan, M. and Li, X. (2015). Does the Green Great Wall effectively decrease dust storm intensity in China? A study based on NOAA NDVI and weather station data. *Land Use Policy*, 43, 42–47. https://www.researchgate.net/publication/268692474_Does_the_Green_Great_Wall_effectively_decrease_dust_storm_intensity_in_China_A_study_based_on_NOAA_NDVI_and_weather_station_data
38. Viña, A., McConnell, W.J., Yang, H., Xu, Z. and Liu, J. (2016). Effects of conservation policy on China's forest recovery. *Science advances*, 2(3), e1500965. <http://advances.sciencemag.org/content/2/3/e1500965.full>
39. UNEP (2015). Review of the Kubuqi Ecological Restoration Project: A Desert Green Economy Pilot Initiative. United Nations Environment Programme, Nairobi. http://wedocs.unep.org/bitstream/handle/20.500.11822/8652/-Review_of_the_Kubuqi_Ecological_Restoration_Project_A_Desert_Green_Economy_Pilot_Initiative-2015Review_of_the_Kubuqi_Ecological_Restoration_Project..pdf?sequence=2&isAllowed=y

40. UNCCD (2017). Great Green Wall. United Nations Convention to Combat Desertification Secretariat, Bonn. <http://www.greatgreenwall.org/great-green-wall/>
41. Sacande, M. and Berrahmouni, N. (2016). Community participation and ecological criteria for selecting species and restoring natural capital with native species in the Sahel. *Restoration Ecology*, 24(4), 479-488. <http://onlinelibrary.wiley.com/doi/10.1111/rec.12337/abstract>
42. UNCCD (2005). *A Master Plan for Regional Cooperation for the Prevention and Control of Dust and Sandstorms*. The Regional Master Plan for the Prevention and Control of Dust and Sandstorms in North East Asia Volume 1. United Nations Convention to Combat Desertification Secretariat, Bonn. http://www.unccd.int/Lists/SiteDocumentLibrary/Publications/dustsandstorms_northeastasia.pdf
43. WMO (2015). *Sand and Dust Storm Warning Advisory and Assessment System (SDS-WAS): Science and Implementation Plan 2015–2020*. World Weather Research Programme Report 2015-5. World Meteorological Organization, Geneva. https://www.wmo.int/pages/prog/arep/wwrp/new/documents/Final_WWRP_2015_5_SDS_IP.pdf
44. WMO (2017). Sand and Dust Storm Warnings website. World Meteorological Organization, Geneva. <https://public.wmo.int/en/our-mandate/focus-areas/environment/sand-and-dust-storm/sand-and-dust-storm-warnings>
45. WOCAT SLM (2017). The Global Database on Sustainable Land Management of the World Overview of Conservation Approaches and Technologies website. University of Bern, Berne. <https://qcat.wocat.net/en/wocat/>
46. FAO (2016). AQUASTAT website. Food and Agriculture Organization of the United Nations, Rome. http://www.fao.org/nr/water/aquastat/water_use/index.stm
47. 2030 WRG (2015). The 2030 Water Resources Group website. <https://www.waterscarcitysolutions.org/#>

Graphic references

48. WHO (2006). *WHO Air quality guidelines for particulate matter, ozone, nitrogen dioxide and sulfur dioxide - Global update 2005*. Summary of risk assessment. World Health Organization, Geneva. http://apps.who.int/iris/bitstream/10665/69477/1/WHO_SDE_PHE_OEH_06.02_eng.pdf
49. Financial Tribune (2017). Dust Storms Slam Khuzestan Again. *Financial Tribune*, 29 January 2017. <https://financialtribune.com/articles/environment/58374/dust-storms-slam-khuzestan-again>
50. Tozer, P. and Leys, J. (2013). Dust storms - what do they really cost? *The Rangeland Journal*, 35, 131-142. <http://www.publish.csiro.au/rj/pdf/RJ12085>



51. Jugder, D., Shinoda, M., Sugimoto, N., Matsui, I., Nishikawa, M., Park, S-U., Chun, Y-S. and Park, M-S. (2011). Spatial and temporal variations of dust concentrations in the Gobi Desert of Mongolia. *Global and Planetary Change*, 78, 14-22. https://www.researchgate.net/publication/241100103_Spatial_and_temporal_variations_of_dust_concentrations_in_the_Gobi_Desert_of_Mongolia
52. Miri, A., Ahmadi, H., Ekhtesasi, M.R., Panjehkeh, N. and Ghanbari, A. (2009). Environmental and socio-economic impacts of dust storms in Sistan Region, Iran. *International Journal of Environmental Studies*, 66, 343-355. <http://www.tandfonline.com/doi/abs/10.1080/00207230902720170?journalCode=genv20>

A dust plume blowing from western Africa towards the Amazon Basin and the Gulf of Mexico, 25 June 2014. At least 40 million tonnes of Saharan dust is delivered to the Amazon Basin each year. ►

Photo credit: Norman Kuring/NASA OceanColor Group